

SAGES


Society of American Gastrointestinal and Endoscopic Surgeons

<http://www.sages.org>

Laparoscopic Spleen Removal (Splenectomy) Patient Information from SAGES

Author : SAGES Webmaster





LAPAROSCOPIC SPLEEN REMOVAL

patient information from your surgeon & SAGES

About the Spleen

What is the Spleen?



The spleen is a blood filled organ located in the upper left abdominal cavity. It is a storage organ for red blood cells and contains many specialized white blood cells called “macrophages” (disease fighting cells) which act to filter blood. The spleen is part of the immune system and also removes old and damaged blood particles from your system. The spleen helps the body identify and kill bacteria. The spleen can affect the platelet count, the red blood cell count and even the white blood count.

How Do I Know If My Spleen Should Be Removed?

There are several reasons why a spleen might need to be removed, and the following list, though not all inclusive, includes the most common reasons.

Auto-immune thrombocytopenia purpura (ITP): This is the most common reason. In this disease, a patient’s platelet count is low because the body makes antibodies to the platelets which cause them to be destroyed in the spleen. Since platelets are blood cells which aid in blood clotting, patients are at risk for bleeding. The first line of treatment is medical therapy but when this is not successful, removal of the spleen (splenectomy) can be curative or at least significantly improve the platelet count in the vast majority of cases.

Hemolytic anemia: In this disease the body makes antibodies to red blood cells which are subsequently destroyed in the spleen. When medical therapy fails, some patients require spleen removal to prevent or decrease the need for transfusion.

Hereditary (genetic) conditions: There are several diseases that affect the shape of red blood cells, such as, spherocytosis, sickle cell disease or thalassemia. In these patients, the spleen recognizes the red cells as abnormal and may bring them down, possibly requiring splenectomy to improve the symptoms.

Malignancy: Rarely, patients with cancers of the cells which fight infection, known as lymphoma or certain types of leukemia, require spleen removal. When the spleen gets enlarged, it sometimes removes too many platelets from your blood and has to be removed. Sometimes the spleen is removed to diagnose or treat a tumor.

Other reasons: Sometimes the blood supply to the spleen becomes blocked (infarct) or the artery abnormally expands (aneurysm) and the spleen needs to be removed. Rarely, the spleen can become infected and is best treated with removal.

About Spleen Removal

How are These Problems Found?

An evaluation typically includes a complete blood count (CBC), to look at the amount, sizes and shapes of the cells in your blood. Sometimes it is important to get a specimen of bone marrow. The bone marrow is where red cells and other components of the blood are made and can be very helpful to understand the source of the problems. Finally, almost nobody has their spleen removed electively without some imaging such as an ultrasound, a computerized tomography (CAT scan), or magnetic resonance imaging (MRI). Understanding the size and shape of the spleen is very helpful to understanding the cause of the problem as well as for surgical planning. In addition, sometimes a nuclear scan of the spleen can be helpful especially if little satellites of the spleen known as accessory spleens or “splenules” are suspected.

What are the Advantages of Laparoscopic Splenectomy?

Individual results may vary depending on your overall condition and health but the usual advantages are:

- Less postoperative pain
- Shorter hospital stay
- Faster return to a regular, solid food diet

- Quicker return to normal activities
- Better cosmetic results
- Fewer [incisional hernias](#)

Am I a Candidate for Laparoscopic Spleen Removal?

Most patients can have a laparoscopic splenectomy. Though the experience of the surgeon is the biggest factor in a successful outcome, the size of the spleen is the most important determinant in deciding whether the spleen can be removed laparoscopically. When the size of the spleen is extremely large, it is difficult to perform the laparoscopic technique. Sometimes, plugging the artery to the spleen right before surgery using special X-ray technology can shrink the spleen to allow the laparoscopic technique. You should obtain a thorough evaluation by a surgeon qualified in laparoscopic spleen removal along with consultation with your other physicians to find out if this technique is appropriate for you.


What Preparation is Required?

- After your surgeon reviews with you the potential risks and benefits of the operation, you will need to provide written consent for surgery.
- Preoperative preparation includes blood work, medical evaluation, chest x-ray and an EKG depending on your age and medical condition.
- Immunization with a vaccine to help prevent bacterial infections after the spleen is removed should be given two weeks before surgery, if possible.
- Blood transfusion and/or blood products such as platelets may be needed depending on your condition.
- It is recommended that you shower the night before or morning of the operation.
- After midnight the night before the operation, you should not eat or drink anything except medications that your surgeon has told you are permissible to take with a sip of water the morning of surgery.
- Drugs such as aspirin, blood thinners, anti-inflammatory medications (arthritis medications) and Vitamin E may need to be stopped temporarily for several days to a week prior to surgery.
- Diet medication or St. John's Wort should not be used for the two weeks prior to surgery.
- Quit smoking and arrange for any help you may need at home.

What should I expect during surgery?

How is Laparoscopic Removal of the Spleen Done?

You will be placed under general anesthesia and be completely asleep. A cannula (hollow tube) is placed into the abdomen by your surgeon and your abdomen will be inflated with carbon dioxide gas to create a space to operate. A laparoscope (a tiny telescope connected to a video camera) is put through one of the cannulas which projects a video picture of the internal organs and spleen on a television monitor. Several cannulas are placed in different locations on your abdomen to allow your surgeon to place instruments inside your belly to work and remove your spleen. A search for accessory (additional) spleens and then removal of these extra spleens will be done since 15% of people have small, extra spleens. After the spleen is cut from all that it is connected to, it is placed inside a special bag. The bag with the spleen inside is pulled up into one of the small, but largest incisions on your abdomen. The spleen is broken up into small pieces (morcelated) within the special bag and completely removed.



What Happens If the Operation Cannot Be Performed or Completed by the Laparoscopic Method?

In a small number of patients the laparoscopic method cannot be performed. Factors that may increase the possibility of choosing or converting to the “open” procedure may include obesity, a history of prior abdominal surgery causing dense scar tissue, inability to visualize organs or bleeding problems during the operation.

The decision to perform the open procedure is a judgment decision made by your surgeon either

before or during the actual operation. When the surgeon feels that it is safest to convert the laparoscopic procedure to an open one, this is not a complication, but rather sound surgical judgment. The decision to convert to an open procedure is strictly based on patient safety.

Expected Outcomes

What Can I Expect after Surgery?

After surgery you will be given intravenous fluids (IV's) in your arm. You will be given pain medication to relieve the discomfort you may experience from the small incisions. You will need to let your nurse and surgeon know what your pain medication needs are since everyone has a different pain threshold.

As soon as you can resume oral intake, urinate, and care for your basic needs, you will typically be able to go home. Your surgeon will tell you when it is safe to go home.

At Home

Typically, once you have gone home, you may do the following, but each situation differs and “at home” activities should be discussed with your doctor.

- **Activity:** You can walk and go up stairs. You can shower, but most surgeons prefer that you do not soak in a tub for at least a week or more after surgery. When you feel stronger ask your surgeon about more strenuous activities.
- **Driving:** Ask your surgeon. Most people can resume driving 5 to 7 days after surgery. You should not be taking pain medication when you drive.
- **Diet:** Unless you have special dietary needs, such as diabetes, you can eat a normal diet at home.
- **Bowel function:** It is common to get constipated after surgery, especially when you are taking pain medication. It is important to drink plenty of water and take in enough fiber in your diet. Your surgeon may advise you about means to help your bowel function postoperatively. Ask or call your surgeon before taking a cathartic.

What Complications Can Occur?

Complications following laparoscopic splenectomy are infrequent, but you should consult your doctor regarding possible complications based on your specific case. Possible complications may include cannula site infections, pneumonia, internal bleeding or infection inside the abdomen at the

site where the spleen used to be, although these complications are infrequent. The pancreas can become inflamed (pancreatitis) or leak fluid after surgery. Problems that can occur a few months to years later are hernias at the cannula sites or overwhelming infection throughout the entire body. This complication is also infrequent.

Overwhelming infection that occurs after splenectomy is called OPSI or Overwhelming Post-Splenectomy Infection. OPSI is a result of not having a spleen to fight certain bacterial infections. Immunization is usually given before you have your spleen removed and is one method to help the body fight and prevent infection. Antibiotics, like penicillin, can be given if an infection develops because the bacteria that commonly cause this type of infection are very sensitive to antibiotics. It is important that you tell your physician or any physician that is covering for your doctor that you had your spleen removed.

When to Call Your Doctor

Be sure to call your physician or surgeon if you develop any of the following:

- Persistent fever over 101 degrees F (39 C)
- Bleeding
- Increasing abdominal swelling
- Pain that is not relieved by your medications
- Persistent nausea or vomiting
- Chills
- Persistent cough or shortness of breath
- Purulent drainage (pus) from any incision
- Redness surrounding any of your incisions that is worsening or getting bigger
- You are unable to eat or drink liquids

This brochure is intended to provide a general overview of a laparoscopic splenectomy. It is not intended to serve as a substitute for professional medical care or a discussion between you and your surgeon about the need for a laparoscopic spleen removal. Specific recommendations may vary among health care professionals. If you have a question about your need for a laparoscopic splenectomy, your alternatives, billing or insurance coverage, or your surgeons training and experience, do not hesitate to ask your surgeon or his/her office staff about it. If you have questions about the operation or subsequent follow up, discuss them with your surgeon before or after the operation.